

Fungicide Efficacy for Control of Wheat Diseases (Revised 3-30-16)

The multi-state North Central Regional Committee on Management of Small Grain Diseases (NCERA-184) has developed the following information on fungicide efficacy for control of certain foliar diseases of wheat in the U.S. Efficacy ratings for each fungicide listed were determined by field testing over multiple years and locations. Efficacy is determined by direct comparisons among products in field tests. The products are used in a manner consistent with label instructions. This table includes most widely marketed products. It is not intended to be a list of all labeled fungicide products.

Fungicides			Pow-dery milde w	Stag. leaf blotch	Septori aleaf spot	Tan spot	Stripe rust	Leaf rust	Stem rust	Head scab	Harvest Restriction	
Active ingredient	Product	Rate oz/ac										
Strobilurin	Picoxystrobin 22.5%	Aproach SC	6.0 – 12.0	G ¹	VG	VG ²	VG	E ³	VG	VG	NL	Feekes 10.5
	Fluoxastrobin 40.3%	Evito 480 SC	2.0 – 4.0	G	--	--	VG	--	VG	--	NL	Feekes 10.5 & 40 days
	Pyraclostrobin 23.6%	Headline SC	6.0 - 9.0	G	VG ²	VG ²	E	E ³	E	G	NL	Feekes 10.5
Triazole	Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	VG	VG	--	VG	E	E	E	G	30 days
	Propiconazole 41.8%	Tilt 3.6 EC ⁴	4.0	VG	VG	VG	VG	VG	VG	VG	P	Feekes 10.5
	Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	--	VG	VG	VG	VG	VG	VG	G	30 days
	Tebuconazole 38.7%	Folicur 3.6 F ⁴	4.0	NL	NL	NL	NL	E	E	E	F	30 days
	Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.2	G	VG	VG	VG	VG	E	E	E	G
Mixed modes of action ⁵	Tebuconazole 22.6% Trifloxystrobin 22.6%	Absolute Maxx SC	5.0	G	VG	VG	VG	VG	E	VG	NL	35 days
	Fluoxastrobin 14.8% Flutriafol 19.3%	Fortix	4.0 - 6.0	--	--	VG	VG	E	VG	--	NL	Feekes 10.5 & 40 days
	Benzovindiflupyr 10.3% Propiconazole 11.7% Azoxystrobin 13.5%	Trivapro A EC + Trivapro B SE	4.0 + 10.5	VG	VG	VG	VG	E	E	VG	NL	?
	Metconazole 7.4% Pyraclostrobin 12%	TwinLine 1.75 EC	7.0 – 9.0	G	VG	VG	E	E	E	VG	NL	Feekes 10.5
	Fluxapyroxad 14.3% Pyraclostrobin 28.6%	Priaxor	4.0 - 8.0	G	VG	VG	E	VG	VG	G	NL	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 13.5%	Quilt Xcel 2.2 SE ⁴	10.5 - 14.0	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5
	Prothioconazole 10.8% Trifloxystrobin 32.3%	Stratego YLD	4.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	Cyproconazole 7.17% Picoxystrobin 17.94%	Aproach Prima SC	3.4 - 6.8	VG	VG	VG	VG	E	VG	--	NR	45 days

¹ Efficacy categories: NL=Not Labeled; NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; "--" connotes Insufficient data to make statement about efficacy.

² Product efficacy may be reduced in areas with fungal populations that are resistant to strobilurin fungicides.

³ Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred.

⁴ Multiple generic products containing the same active ingredients also may be labeled in some states.

For more information, contact:
 Martin Nagelkirk nagelkir@msu.edu
 Martin Chilvers chilvers@msu.edu

⁵ *Products with mixed modes of action generally combine triazole and strobilurin active ingredients.*

However, Priaxor and the Trivapro co-pack also include carboxamide active ingredients.